

Asparagine synthetase: Function, structure, and role in disease

Carrie L. Lomelino^{1,†}, Jacob T. Andring^{1,†}, Robert McKenna^{1,‡}, Michael S. Kilberg^{1,‡}

¹Department of Biochemistry and Molecular Biology, Shands Cancer Center, College of Medicine, University of Florida, Gainesville, Florida 32610, USA.

† Co-first authors

‡ Co-senior authors

Running title: Asparagine Synthetase: Role in Disease

To whom correspondence should be addressed: Michael S. Kilberg, Department of Biochemistry and Molecular Biology, College of Medicine, University of Florida, Gainesville, Florida 32610, USA. Telephone: 352-294-8388, email: mkilberg@ufl.edu

Abstract:

Asparagine synthetase (ASNS) converts aspartate and glutamine to asparagine and glutamate in an ATP-dependent reaction. ASNS is present in most, if not all, mammalian organs, but varies widely in basal expression. Human ASNS activity is highly responsive to cellular stress, primarily by increased transcription from a single gene located on chromosome 7. Elevated ASNS protein expression is associated with resistance to asparaginase therapy in childhood acute lymphoblastic leukemia. There is evidence that ASNS expression levels may also be inversely correlated with asparaginase efficacy in certain solid tumors as well. Children with mutations in the ASNS gene exhibit developmental delays, intellectual disability, microcephaly, intractable seizures, and progressive brain atrophy. Thus far, 15 unique mutations in the ASNS gene have been clinically associated with asparagine synthetase deficiency (ASD). Molecular modelling using the *E. coli* ASNS-B structure has revealed that most of the reported ASD substitutions are located near catalytic sites or within highly conserved regions of the protein. For some ASD patients, fibroblast cell culture studies have eliminated protein and mRNA synthesis or stability as the basis for decreased proliferation.

ASNS Function

Asparagine synthetase (ASNS) catalyzes the synthesis of asparagine and glutamate from aspartate and glutamine in an ATP-dependent amidotransferase reaction (Fig. 1A) (1). The recent discovery of a neurologic disorder associated with asparagine synthetase deficiency (ASD) and its long recognized importance in the treatment of acute lymphoblastic leukemia (ALL) highlight the clinical relevance of ASNS as a topic of current interest.

Many prokaryotes express two forms of ASNS that are characterized by their source of nitrogen donor, either ammonia (AS-A) or glutamine (AS-B) (1). The highly conserved residue Cys2 is required for the nucleophilic deamidation of glutamine and the mutation of this residue reverts AS-B to use ammonia exclusively as the nitrogen source (2). In mammalian cells, a single form of the ASNS enzyme is expressed that utilizes glutamine as the nitrogen donor in a reaction corresponding to bacterial AS-B (2). Structure-function studies have previously been performed on *E. coli* AS-B and have revealed two distinct catalytic domains that are conserved in the human enzyme (3) (Fig. 1B). Enzymatic analyses have shown that the catalytic mechanism requires magnesium ions and ATP (4). The reaction begins in the C-terminal domain with the activation of the aspartate carboxyl group. This ATP-dependent process forms a β -aspartyl-AMP intermediate that is stabilized by proximal residues of the active site. Within the N-terminal domain, the hydrolysis of glutamine yields glutamate and ammonia, the latter of which diffuses through an intramolecular tunnel to perform a nucleophilic attack on the electrophilic β -aspartyl-AMP intermediate, producing asparagine (Fig. 1A) (1,3,5). Human ASNS is categorized as a class II or Ntn glutamine amidotransferase because the hydrolysis of glutamine occurs in the N-terminal domain of the enzyme (1).

ASNS protein Structure

The crystal structure of *E. coli* AS-B (PDB: 1CT9) has been solved (3), but a structure of human ASNS has yet to be published or deposited into the protein database (PDB). Therefore, the structural features of the human enzyme are based on homology modelling using the *E. coli* AS-B structure as a template because they share approximately 40% sequence identity. The canonical human ASNS enzyme consists of 561 amino acid residues with an approximate molecular mass of 65 kDa (Fig. 1B). However, the UniProt protein database contains two other putative isoforms of

human ASNS that differ in the length of the N terminal domain: isoform 2 (residues 22-561) and isoform 3 (residues 84-561). The expression and physiological importance of these putative isoforms have not been described. With regard to the two functional domains, the N-terminal domain (residues 1-208) consists of a two layer, antiparallel β -sheet core surrounded by four α -helices (Fig. 1C). This domain contains the glutamine binding pocket, consisting of residues R49, N75, E77, and D97. As observed in the *E. coli* structure, glutamine is predicted to bind in a manner so that the carboxamide group is oriented toward the interface of the two domains to allow the transfer of an ammonia group from glutamine to aspartate. The C-terminal domain (residues 209-561) is composed primarily of α -helices, but also encompasses a 5-stranded, parallel β -sheet that contains the ATP binding site: residues L256, V288, D295, S363, G364, E365, and D401 (Fig. 1C). A distance of approximately 20Å separates the two active sites.

Asparagine Synthetase and Cancer

ASNS expression and asparagine metabolism have received considerable attention in transformed cells, beginning with the observation that childhood ALL is susceptible to treatment by the infusion of bacterial asparaginase (ASNase). Primary ALL cells and many ALL cell lines exhibit little or no detectable ASNS and are sensitive to asparagine depletion (6-8). Standard treatment of childhood ALL includes infusion of bacterial ASNase as a principle component of a combinatorial chemotherapy (9). The circulating ASNase causes rapid depletion of plasma asparagine, followed by the efflux of intracellular asparagine and thus, starving the leukemia cells to prevent further growth (10,11). In contrast to the increase in ASNS expression in response to substrate deprivation in most normal cells within the body, there is little or no up-regulation of ASNS protein content in ALL cells rendering them preferentially sensitive to ASNase (12,13).

The importance of ASNS expression in solid tumor growth and the sensitivity to ASNase has not been investigated as extensively. A screen of 98 human pancreatic ductal carcinomas established that ASNS protein was low or below detection in about 70% of the samples, suggesting that some pancreatic tumors may be susceptible to ASNase therapy (14). Cui et al. showed that pancreatic cancer cells over-expressing ASNS exhibited increased resistance to apoptosis induced by cisplatin (15). Similarly, using the NCI-60 human cancer cell panel, Lorenzi et al. noted a negative correlation between ASNS mRNA levels and susceptibility to ASNase in ovarian cells, as well as increased ASNase sensitivity after siRNA knockdown of ASNS expression (16). In a second study with a larger number of ovarian cell lines, an inverse correlation between ASNase efficacy and ASNS protein levels was observed (17). These results were consistent with previous observations in human leukemia cells showing that increased ASNase sensitivity correlated with lower ASNS protein expression rather than mRNA content (13).

Protein limitation or an imbalanced dietary amino acid composition lead to intracellular amino acid depletion and activation of ASNS gene transcription through a signaling pathway called the amino acid response (AAR) (18). Likewise, endoplasmic reticulum stress also increases ASNS transcription via the unfolded protein response (UPR) (19). These cell stress pathways result in activation of the eIF2 kinases GCN2 (AAR) and PERK (UPR). Phosphorylation of the alpha subunit of eIF2 causes a transient, partial suppression of global protein synthesis, but a paradoxical increase in the translation for certain mRNAs and among these is the transcription factor ATF4 (20). ATF4 binds to an enhancer element within the proximal promoter of the ASNS gene and activates transcription (Fig. 2) (21). Ye et al. investigated the role of ATF4 in tumor cell survival and proliferation and observed that ATF4 knockdown caused reduced survival in HT1080 fibrosarcoma and

DLD1 colorectal adenocarcinoma cells in the absence of non-essential AA (22). Reduced proliferative capacity and increased apoptosis correlated with lower ASNS expression in the ATF4-deficient cells. Supplementation of the tumor cells with asparagine, but not other amino acids, led to increased cell survival. Based on this and other experimental data, Ye et al. concluded that induction of ATF4: 1) contributes to tumor cell proliferation during nutrient limitation, 2) is a component of starvation-induced autophagy in cancer cells, and 3) induces ASNS as a key factor in tumor initiation and growth under AA limiting conditions (22). The exact role of ASNS activity in modulating tumor growth is unknown. One obvious possibility, an impact on protein synthesis, seems tenuous as other amino acid synthetic enzymes are not as highly regulated as ASNS.

Asparagine Synthetase Deficiency

Asparagine Synthetase Deficiency (ASD) is a recently characterized neurological disorder having severe impacts on psychomotor development and mortality at an early age. Multiple patient studies have been conducted over the last few years due to the rising awareness of the disease. Symptoms include intellectual disability, microcephaly, severe developmental delay, intractable seizures, progressive brain atrophy, and more rarely, respiratory deficiency (23-30). The disease is associated with homozygous or compound heterozygous mutations within the ASNS gene on chromosome 7q2, but the exact mechanisms that cause the overt symptoms of the disease are not well understood (23,26,31). The fact that the children are born with epileptic-like seizures and microcephaly suggests that brain ASNS activity is critical for development of this organ. Currently, the disease can only be diagnosed through DNA sequencing. Some, but not all, affected individuals have a measurable decrease in the amount of asparagine in their serum or cerebrospinal fluid, which limits this analysis as a preliminary screen in suspected cases.

Based on animal growth studies, asparagine is traditionally defined as a “non-essential” or “indispensable” amino acid. ASNS is expressed to varying degrees within human organs (<http://www.proteinatlas.org/ENSG00000070669-ASNS/tissue>), but particularly high in pancreas (32). However, when considering the requirement for asparagine at the level of individual cells, ASNS deficiency leads to extracellular asparagine dependence, as discussed above for ALL cells. Individuals with ASD express phenotypic impairments in the brain that are not readily apparent in other organs, suggesting a tissue-specific dependence on asparagine for neural development (23). Although asparagine can be obtained through the diet, it is transported by an equilibrating, bi-directional facilitated transporter (System n) across the luminal surface of the endothelial cells that make up the blood brain barrier. Therefore, asparagine is not actively accumulated within the brain (33). Indeed, Na⁺-dependent transport of asparagine across the abluminal membrane of the endothelial layer appears to be designed to remove the amino acid from the brain. Consistent with this hypothesis, the CSF levels of asparagine, as with many amino acids, are only a fraction of those in the plasma (34,35). Consequently, a decrease in ASNS catalytic activity in the brain is presumed to cause the disease phenotype (23,24,26). Plasma asparagine levels were reduced in 8/16 ASD patients for whom plasma analysis was reported (reviewed in 31). However, CSF asparagine content has been analyzed in only four ASD patients and it was undetectable in two of the four (24,27,30). Recent studies with ASD patients' fibroblasts have revealed that lowering the asparagine concentration in the extracellular milieu results in an inability to proliferate (26). These cell culture studies also showed that fibroblasts from asymptomatic heterozygote parents exhibit no ill effects in the presence of sufficient extracellular asparagine, but decline in proliferative capacity when placed in an asparagine-free medium. Collectively,

investigation of ASD patients indicates that ASNS activity in the brain is crucial for organ development. Whether this dependence is the direct result of perturbations in asparagine metabolism, or one of the other ASNS reactants must be established by further analysis and experimentation.

Structural Examination of ASD Mutations

To date, 15 unique mutations in the ASNS gene have been clinically identified in association with ASD (23-31). ASD is characterized as a pan-ethnic disorder because it has been identified within a variety of ethnic groups. Several patients have mutations localized at or near the ATP binding site, whereas thus far, only one patient has been identified with a mutation in the glutamine binding site (Table 1). The remaining mutations are located outside the active sites, but in highly conserved regions, that may reduce protein stability without necessarily inhibiting enzyme activity (23). Alignment of the three isoforms of human ASNS and renumbering of the mutations to correlate to the isoform 1 sequence highlights that previously reported mutations Y315C (27) and Y377C (24) are equivalent to Y398C (25), whereas R324* would be renumbered to R407* (27) (Fig. 1B).

Cell culture analyses have shown that several of the aforementioned mutations do not affect mRNA stability (23,26). Rather, many of the disease-associated mutations are expected to decrease enzyme activity and/or protein stability. Analysis of the human ASNS structural model incorporating the observed amino acid mutations illustrates an overall theme of protein destabilization. The mutations can be classified into four groups based upon their putative effect on the protein structure: change in the amino acid type (hydrophobic to hydrophilic or *vice versa*), loss of hydrogen bonding or van der Waals interactions, truncation of the protein, and predicted decrease in substrate binding. The introduction of charged or polar residues into hydrophobic regions of the enzyme is expected to destabilize the structure as predicted for the A6E, L145S, A380S, and

V489D mutations (23,30). In addition, the increase in side chain length of the A6E mutation results in steric hindrance with proximal residue F8. Conversely, the S480F mutation presents a nonpolar amino acid on the surface of protein that may decrease solubility (Fig. 3) (28). Several mutations result in the loss of stabilizing interactions. L247W causes a reorientation of the larger aromatic side chain out of a hydrophobic pocket to avoid steric clash, resulting in the loss of van der Waals interactions (30). Similarly, the reduction in side chain size is likely to cause a decrease in van der Waals interactions in F362V (23) and a loss of hydrogen bonding in the R340H mutant (29), which is further destabilized by steric hindrance with F482. In addition to the introduction of a reactive, solvent accessible thiol group, the loss of van der Waals interactions in Y398C is predicted to disrupt contacts between the N and C domains (24,25,27). Although the R550C mutation is not included in the human ASNS model due to the lack of inclusion in the *E. coli* AS-B structure, it is expected to result in destabilization of the C-terminal domain (Fig. 3).

Two observed mutations are predicted to result in the truncation of the human ASNS protein. The nonsense mutation R407* introduces a premature stop codon (27), whereas the frame shift mutation W541Cfs*5 truncates the enzyme, containing only five codons after the mutated residue 541 (30). The remaining reported mutations are not only predicted to destabilize the structure of the protein, but also potentially decrease the binding of substrate and possibly decrease the catalytic efficiency of the ASNS enzyme. R49Q is the first mutation reported to be in the glutamine binding pocket of the N-terminal domain (publication submitted for review, Sacharow, Dudenhausen, Lomelino, McKenna, Kilberg, and Rodan). This mutation not only results in a loss of hydrogen bonds with the second beta sheet, but also a loss in hydrogen bonds with the glutamine substrate. The C-terminal domain mutations G289A and T337I are associated with an afflicted child

who is a compound heterozygote. Both mutations are both located proximal to the ATP binding pocket (26). The G289A mutation results in a steric clash with Ser293, whereas the T337I mutation introduces a hydrophobic patch on the protein surface that may decrease solubility. Although these two residues are not predicted to interact directly with the ATP molecule, the mutations are expected to destabilize the region of the ATP binding site and therefore, potentially decrease the affinity of ASNS for this substrate (Fig. 3).

Remaining Questions

Although many aspects of ASNS enzymology and regulation have been investigated over the last four decades, major gaps remain in our knowledge. The exact contribution of ASNS activity in maintaining the cellular and whole body homeostasis of its four amino acid reactants remains largely unanswered. Although the enzyme name leads one to focus on asparagine synthesis, given the overall reaction, the activity does consume glutamine and aspartate, and produce glutamate. Whether or not ASNS actually impacts the cellular homeostasis of one or more of the other reactants speculative. Given the much higher levels of glutamine in most cells, the chance of changes in ASNS activity contributing to the cellular aspartate and glutamate abundance would be more likely. However, the *in vivo* impact during embryonic development or physiological status during periods of up-regulation in response to cellular stresses that activate the AAR or UPR pathways would be valuable studies in tissue-specific ASNS knockout animals. Furthermore, although limited, the results showing the importance of increased ASNS expression in solid tumor proliferation and the development of resistance to ASNase chemotherapy in childhood ALL illustrate the need for a better comprehension of ASNS regulation in transformed cells.

The recent awareness of the neurological disease ASD highlights the need

for a better understanding of the enzymatic consequences of ASNS mutations. The pioneering mouse model studies of Ruzzo et al. should be expanded to generate brain-specific transgenic and knockout animals to investigate the developmental effects of altered ASNS activity (23). Given that fibroblasts are available from only a few ASD children, the physiologic consequences for some mutations will require expression in ASNS null cells. At the protein level, modeling of the human ASNS protein has provided a means to tentatively map the structural consequences of ASD-associated mutations. However, the x-ray crystal structure of the human enzyme would be useful to confirm the structural effects proposed for ASD and allow for potential *in silico* drug screening approaches for future anti-tumor use. Furthermore, the current enzymatic assays for ASNS are cumbersome and variable in reproducibility when performed at the relatively low tissue levels of the enzyme. Over-expression of wild type and mutant recombinant human ASNS protein in quantities suitable for new attempts at analysis would allow for activity comparisons of each mutation. With regard to ASD, given that asparagine uptake across the blood brain barrier is not concentrative may limit dietary amino acid supplement therapy because artificially elevating blood asparagine levels too high may inhibit uptake of other amino acids due to competition for shared transporters (33). In fact, an initial attempt at dietary asparagine therapy resulted in negative consequences (36). Affected ASD children are born with morphological and neurologic defects, indicating significant brain damage during embryonic development, and ASNS has been proven essential during brain development in a mouse model (23). These observations suggest that currently available therapeutic approaches are likely to prove ineffective, as significant and possibly irreversible tissue damage appears to occur during the earliest stages of development.

Conflict of Interest

The authors declare that they have no conflicts of interest with the contents of this article.

Author contributions

RM conceived the report, CLL, JTA, and RM reviewed the literature, wrote the text, and prepared the figures describing the ASNS structure-function studies and ASD. MSK contributed the sections on regulation of the ASNS gene, ASD, and ALL chemotherapy. All authors worked together to revise and improve each successive version of the manuscript.

References

1. Richards, N. G., and Kilberg, M. S. (2006) Asparagine synthetase chemotherapy. *Annu. Rev. Biochem.* **75**, 629-654
2. Richards, N. G. J., and Schuster, S. M. (1998) Mechanistic issues in asparagine synthetase catalysis. *Adv. Enzymol.* **72**, 145-198
3. Larsen, T. M., Boehlein, S. K., and Schuster, S. M. (1999) Three-dimensional structure of *Escherichia coli* asparagine synthetase B: A short journey from substrate to product. *Biochemistry (Mosc)*. **38**, 16146-16157
4. Patterson, M. K., Jr., and Orr, G. R. (1968) Asparagine biosynthesis by the Novikoff Hepatoma isolation, purification, property, and mechanism studies of the enzyme system. *J. Biol. Chem.* **243**, 376-380
5. Tesson, A. R., Soper, T. S., Ciustea, M., and Richards, N. G. (2003) Revisiting the steady state kinetic mechanism of glutamine-dependent asparagine synthetase from *Escherichia coli*. *Arch. Biochem. Biophys.* **413**, 23-31
6. Broome, J. D. (1968) Studies on the mechanism of tumor inhibition by L-asparaginase. *J. Exp. Med.* **127**, 1055-1072
7. Asselin, B. L., and Kurtzburg, J. (2003) Asparaginase. in *Treatment of Acute Leukemias* (Pui, C. H. ed.), Humana Press, Totawa, NJ. pp 365-379
8. Rizzari, C. (2003) Asparaginase Treatment. in *Treatment of Acute Leukemias* (Pui, C. H. ed.), Humana Press, Towata, NJ. pp 381-391
9. Pieters, R., Hunger, S. P., Boos, J., Rizzari, C., Silverman, L., Baruchel, A., Goekbuget, N., Schrappe, M., and Pui, C. H. (2011) L-asparaginase treatment in acute lymphoblastic leukemia: a focus on Erwinia asparaginase. *Cancer* **117**, 238-249
10. Aslanian, A. M., and Kilberg, M. S. (2001) Multiple adaptive mechanisms affect asparagine synthetase substrate availability in asparaginase-resistant MOLT-4 human leukaemia cells. *Biochem. J.* **358**, 59-67
11. Avramis, V. I. (2012) Asparaginases: biochemical pharmacology and modes of drug resistance. *Anticancer Res.* **32**, 2423-2437
12. Aslanian, A. M., Fletcher, B. S., and Kilberg, M. S. (2001) Asparagine synthetase expression alone is sufficient to induce L-asparaginase resistance in MOLT-4 human leukaemia cells. *Biochem. J.* **357**, 321-328
13. Su, N., Pan, Y. X., Zhou, M., Harvey, R. C., Hunger, S. P., and Kilberg, M. S. (2008) Correlation between asparaginase sensitivity and asparagine synthetase protein content, but not mRNA, in acute lymphoblastic leukemia cell lines. *Pediatr. Blood Cancer* **50**, 274-279
14. Dufour, E., Gay, F., Aguera, K., Scoazec, J. Y., Horand, F., Lorenzi, P. L., and Godfrin, Y. (2012) Pancreatic tumor sensitivity to plasma L-asparagine starvation. *Pancreas* **41**, 940-948
15. Cui, H., Darmanin, S., Natsuisaka, M., Kondo, T., Asaka, M., Shindoh, M., Higashino, F., Hamuro, J., Okada, F., Kobayashi, M., Nakagawa, K., and Koide, H. (2007) Enhanced Expression of Asparagine Synthetase under Glucose-Deprived Conditions Protects Pancreatic Cancer Cells from Apoptosis Induced by Glucose Deprivation and Cisplatin. *Cancer Res.* **67**, 3345-3355

16. Lorenzi, P. L., Reinhold, W. C., Rudelius, M., Gunsior, M., Shankavaram, U., Bussey, K. J., Scherf, U., Eichler, G. S., Martin, S. E., Chin, K., Gray, J. W., Kohn, E. C., Horak, I. D., Von Hoff, D. D., Raffeld, M., Goldsmith, P. K., Caplen, N. J., and Weinstein, J. N. (2006) Asparagine synthetase as a causal, predictive biomarker for L-asparaginase activity in ovarian cancer cells. *Mol. Cancer Ther.* **5**, 2613-2623
17. Lorenzi, P. L., Llamas, J., Gunsior, M., Ozbun, L., Reinhold, W. C., Varma, S., Ji, H., Kim, H., Hutchinson, A. A., Kohn, E. C., Goldsmith, P. K., Birrer, M. J., and Weinstein, J. N. (2008) Asparagine synthetase is a predictive biomarker of L-asparaginase activity in ovarian cancer cell lines. *Mol. Cancer Ther.* **7**, 3123-3128
18. Kilberg, M. S., Balasubramanian, M., Fu, L., and Shan, J. (2012) The transcription factor network associated with the amino Acid response in Mammalian cells. *Adv Nutr* **3**, 295-306
19. Kilberg, M. S., Shan, J., and Su, N. (2009) ATF4-Dependent Transcription Mediates Signaling of Amino Acid Limitation. *Trends Endocrinol Metab* **20**, 436-443
20. Baird, T. D., and Wek, R. C. (2012) Eukaryotic initiation factor 2 phosphorylation and translational control in metabolism. *Adv Nutr* **3**, 307-321
21. Balasubramanian, M. N., Butterworth, E. A., and Kilberg, M. S. (2013) Asparagine Synthetase: Regulation by Cell Stress and Involvement in Tumor Biology. *American journal of physiology. Endocrinology and metabolism* **304**, E789-799
22. Ye, J., Kumanova, M., Hart, L. S., Sloane, K., Zhang, H., DePanis, D. N., Bobrovnikova-Marjon, E., Diehl, A., Ron, D., and Koumenis, C. (2010) The GCN2-ATF4 pathway is critical for tumor cell survival and proliferation in response to nutrient deprivation. *EMBO J.* **29**, 2082-2096
23. Ruzzo, E. K., Capo-Chichi, J. M., Ben-Zeev, B., Chitayat, D., Mao, H., Pappas, A. L., Hitomi, Y., Lu, Y. F., Yao, X., Hamdan, F. F., Pelak, K., Reznik-Wolf, H., Bar-Joseph, I., Oz-Levi, D., Lev, D., Lerman-Sagie, T., Leshinsky-Silver, E., Anikster, Y., Ben-Asher, E., Olender, T., Colleaux, L., Decarie, J. C., Blaser, S., Banwell, B., Joshi, R. B., He, X. P., Patry, L., Silver, R. J., Dobrzaniecka, S., Islam, M. S., Hasnat, A., Samuels, M. E., Aryal, D. K., Rodriguiz, R. M., Jiang, Y. H., Wetsel, W. C., McNamara, J. O., Rouleau, G. A., Silver, D. L., Lancet, D., Pras, E., Mitchell, G. A., Michaud, J. L., and Goldstein, D. B. (2013) Deficiency of asparagine synthetase causes congenital microcephaly and a progressive form of encephalopathy. *Neuron* **80**, 429-441
24. Alfadhel, M., Alrifai, M. T., Trujillano, D., Alshaalan, H., Al Othaim, A., Al Rasheed, S., Assiri, H., Alqahtani, A. A., Alaamery, M., Rolfs, A., and Eyaid, W. (2015) Asparagine Synthetase Deficiency: New Inborn Errors of Metabolism. *JIMD Rep* **22**, 11-16
25. Ben-Salem, S., Gleeson, J. G., Al-Shamsi, A. M., Islam, B., Hertecant, J., Ali, B. R., and Al-Gazali, L. (2015) Asparagine synthetase deficiency detected by whole exome sequencing causes congenital microcephaly, epileptic encephalopathy and psychomotor delay. *Metab. Brain Dis.* **30**, 687-694
26. Palmer, E. E., Hayner, J., Sachdev, R., Cardamone, M., Kandula, T., Morris, P., Dias, K. R., Tao, J., Miller, D., Zhu, Y., Macintosh, R., Dinger, M. E., Cowley, M. J., Buckley, M. F., Roscioli, T., Bye, A., Kilberg, M. S., and Kirk, E. P. (2015) Asparagine Synthetase Deficiency causes reduced proliferation of cells under conditions of limited asparagine. *Mol. Genet. Metab.* **116**, 178-186
27. Seidahmed, M. Z., Salih, M. A., Abdulbasit, O. B., Samadi, A., Al Hussien, K., Miqdad, A. M., Biary, M. S., Alazami, A. M., Alorainy, I. A., Kabiraj, M. M., Shaheen, R., and Alkuraya,

- F. S. (2016) Hyperekplexia, microcephaly and simplified gyral pattern caused by novel ASNS mutations, case report. *BMC Neurol* **16**, 105-111
28. Gataullina, S., Lauer-Zillhardt, J., Kaminska, A., Galmiche-Rolland, L., Bahi-Buisson, N., Pontoizeau, C., Ottolenghi, C., Dulac, O., and Fallet-Bianco, C. (2016) Epileptic Phenotype of Two Siblings with Asparagine Synthetase Deficiency Mimics Neonatal Pyridoxine-Dependent Epilepsy. *Neuropediatrics* **47**, 399-403
 29. Sun, J., McGillivray, A. J., Pinner, J., Yan, Z., Liu, F., Bratkovic, D., Thompson, E., Wei, X., Jiang, H., Asan, and Chopra, M. (2016) Diaphragmatic Eventration in Sisters with Asparagine Synthetase Deficiency: A Novel Homozygous ASNS Mutation and Expanded Phenotype. *JIMD Rep* **34**, 1-9
 30. Yamamoto, T., Endo, W., Ohnishi, H., Kubota, K., Kawamoto, N., Inui, T., Imamura, A., Takanashi, J. I., Shiina, M., Saitsu, H., Ogata, K., Matsumoto, N., Haginoya, K., and Fukao, T. (2017) The first report of Japanese patients with asparagine synthetase deficiency. *Brain Dev.* **39**, 236-242
 31. Gupta, N., Tewari, V. V., Kumar, M., Langeh, N., Gupta, A., Mishra, P., Kaur, P., Ramprasad, V., Murugan, S., Kumar, R., Jana, M., and Kabra, M. (2017) Asparagine Synthetase deficiency-report of a novel mutation and review of literature. *Metab. Brain Dis.*
 32. Milman, H. A., and Cooney, D. A. (1974) The distribution of L-asparagine synthetase in the principal organs of several mammalian and avian species. *Biochem. J.* **142**, 27-35
 33. Hawkins, R. A., O'Kane, R. L., Simpson, I. A., and Vina, J. R. (2006) Structure of the blood-brain barrier and its role in the transport of amino acids. *J. Nutr.* **136**, 218S-226S
 34. Scholl-Burgi, S., Haberlandt, E., Heinz-Erian, P., Deisenhammer, F., Albrecht, U., Sigl, S. B., Rauchenzauner, M., Ulmer, H., and Karall, D. (2008) Amino acid cerebrospinal fluid/plasma ratios in children: influence of age, gender, and antiepileptic medication. *Pediatrics* **121**, e920-926
 35. Akiyama, T., Kobayashi, K., Higashikage, A., Sato, J., and Yoshinaga, H. (2014) CSF/plasma ratios of amino acids: reference data and transports in children. *Brain Dev.* **36**, 3-9
 36. Alrifai, M. T., and Alfadhel, M. (2016) Worsening of Seizures After Asparagine Supplementation in a Child with Asparagine Synthetase Deficiency. *Pediatr. Neurol.* **58**, 98-100

FOOTNOTES

Acknowledgements - MSK was supported by a grant from the National Cancer Institute (CA-203565). CLL is supported by the National Center for Advancing Translational Sciences of the National Institutes of Health under University of Florida Clinical and Translational Science Awards TL1TR001428 and UL1TR001427.

The abbreviations used are: AA, amino acids; AAR, amino acid response; ALL, acute lymphoblastic leukemia; ASD, asparagine synthetase deficiency; ASNase, asparaginase; ASNS, asparagine synthetase; ATF, activating transcription factor; GCN2, general control non-derepressible; PERK, PKR-like endoplasmic reticulum kinase; UPR, unfolded protein response

Figure Legends

Figure 1. Mechanism and structural features of human asparagine synthetase. A. The reaction begins when the aspartate carboxyl is activated by an ATP-dependent process, forming a β -aspartyl-AMP intermediate. Glutamine deamidation releases ammonia, which performs a nucleophilic attack on the aspartyl intermediate to produce asparagine. Glutamate is the second product of the overall reaction. **B.** Sequence of human ASNS isoform 1 with residues colored light and dark gray for the N- and C-terminal domains, respectively. Alpha helical and beta sheet secondary structures are shown above the sequence. Residues associated with glutamine and ATP binding are colored purple and orange, respectively. Clinically identified ASD mutations are colored according to Fig. 3. **C.** The N and C-terminal domains are represented within the surface structure and colored light and dark gray, respectively. The substrates glutamine (purple) and ATP (orange) are indicated by an arrow and shown as sticks. Insets show the substrate binding pockets. Hydrogen bonds are shown as black dashes with distances labelled in angstroms.

Figure 2. Regulation of ASNS expression. Asparagine depletion activates the amino acid response (AAR), whereas ER stress activates the unfolded protein response (UPR). Each stress condition increases the activity of an eIF2 kinase. Phosphorylation of eIF2 slows global protein synthesis, but paradoxically increases translation of a subset of mRNAs, including that for the transcription factor ATF4. Binding of ATF4 to an enhancer element within the promoter of the ASNS gene induces expression of the enzyme.

Figure 3. ASD associated mutations in the human ASNS enzyme. Mutations are represented as sticks within the predicted ASNS structure. **Top Panel:** A6E (red), L145S (salmon), T337I (light green), R340H (dark green), A380S (light blue), and Y398C (dark blue). **Bottom Panel:** (180° rotation around y-axis) R49Q (raspberry), L247W (light orange), G289A (yellow), F362V (teal), S480F (light pink), V489D (dark pink). Interacting residues are also shown as sticks and hydrogen bonds are represented as black dashes

Table 1. Reported mutations in ASD patients. Residue numbers correspond to those for the isoform 1 ASNS sequence (UniProt ID: P08243).

Mutation	Type	Structural Consequence	Reference
A6E	CH	Charged amino acid in hydrophobic region, steric clash with F8	(23)
R550C		Decrease in side chain length likely to result in loss of interactions	
R49Q	H	GLN binding site, loss of hydrogen bonding	Sacharow et al. (unpublished)
L145S	CH	Polar side chain in hydrophobic region	(30)
L247W		Decrease in van der Waals interactions	
G289A	CH	Proximal to ATP binding site, steric hindrance with Ser293	(26)
T337I		Proximal to ATP binding site, hydrophobic patch on protein that may decrease solubility	
R340H	H	Loss of hydrogen bonds, steric clash with F482	(29)
F362V	H	Decrease in van der Waals interactions	(23)
A380S	H	Polar residue in hydrophobic region	(31)
Y398C	H	Decrease in van der Waals interactions, solvent accessible thiol group	(24,25,27)
R407*	H	Protein truncation	(27)
S480F	CH	Nonpolar residue on protein surface that may decrease solubility	(28)
R550C		Decrease in side chain length likely to result in loss of interactions	
V489D	CH	Charged amino acid in hydrophobic region	(30)
W541Cfs*5		Protein truncation	
R550C	H	Decrease in side chain length likely to result in loss of interactions	(23)

*H= homozygous, CH= compound heterozygous, *=nonsense mutation, fs= frameshift mutation*

Figure 1.




Figure 2.





Figure 3.



Asparagine synthetase: Function, structure, and role in disease
Carrie L Lomelino, Jacob T Andring, Robert McKenna and Michael S. Kilberg

J. Biol. Chem. published online October 30, 2017

Access the most updated version of this article at doi: [10.1074/jbc.R117.819060](https://doi.org/10.1074/jbc.R117.819060)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts